

THE LANCET

Microbe

Supplementary appendix

This appendix formed part of the original submission and has been peer reviewed.
We post it as supplied by the authors.

Supplement to: Chin A W H, Chu J T S, Perera M R A, et al. Stability of SARS-CoV-2 in different environmental conditions. *Lancet Microbe* 2020; published online April 2.
[https://doi.org/10.1016/S2666-5247\(20\)30003-3](https://doi.org/10.1016/S2666-5247(20)30003-3).

Table. Stability of SARS-CoV-2 at different environmental conditions.

A) Temperature*

Time	Virus titre (Log TCID ₅₀ /mL)									
	4°C		22°C		37°C		56°C		70°C	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
1 min	N.D.	N.D.	6.51	0.27	N.D.	N.D.	6.65	0.1	5.34	0.17
5 mins	N.D.	N.D.	6.7	0.15	N.D.	N.D.	4.62	0.44	U	-
10 mins	N.D.	N.D.	6.63	0.07	N.D.	N.D.	3.84	0.32	U	-
30 mins	6.51	0.27	6.52	0.28	6.57	0.17	U	-	U	-
1 hr	6.57	0.32	6.33	0.21	6.76	0.05	U	-	U	-
3 hrs	6.66	0.16	6.68	0.46	6.36	0.19	U	-	U	-
6 hrs	6.67	0.04	6.54	0.32	5.99	0.26	U	-	U	-
12 hrs	6.58	0.21	6.23	0.05	5.28	0.23	U	-	U	-
1 day	6.72	0.13	6.26	0.05	3.23	0.05	U	-	U	-
2 days	6.42	0.37	5.83	0.28	U	-	U	-	U	-
4 days	6.32	0.27	4.99	0.18	U	-	U	-	U	-
7 days	6.65	0.05	3.48	0.24	U	-	U	-	U	-
14 days	6.04	0.18	U	-	U	-	U	-	U	-

B) Surfaces*

Time	Virus titre (Log TCID ₅₀ /ml)									
	Paper		Tissue paper		Wood		Cloth		Glass	
	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD	Mean	±SD
0 min	4.76	0.10	5.48	0.10	5.66	0.39	4.84	0.17	5.83	0.04
30 mins	2.18	0.05	2.19	0.17	3.84	0.39	2.84	0.24	5.81	0.27
3 hrs	U	-	U	-	3.41	0.26	2.21 [#]	-	5.14	0.05
6 hrs	U	-	U	-	2.47	0.23	2.25	0.08	5.06	0.31
1 day	U	-	U	-	2.07 [#]	-	2.07 [#]	-	3.48	0.37
2 days	U	-	U	-	U	-	U	-	2.44	0.19
4 days	U	-	U	-	U	-	U	-	U	-
7 days	U	-	U	-	U	-	U	-	U	-

Time	Banknote					Stainless steel					Plastic					Mask, inner layer			Mask, outer layer		
	Mean		±SD		Mean		±SD		Mean		±SD		Mean		±SD		Mean		±SD		
0 min	6.05	0.34	5.80	0.02	5.81	0.03	5.88	0.69	5.78	0.10	5.83	0.29	5.83	0.04	5.75	0.08	5.75	0.29	5.11	0.51	
30 mins	5.83	0.29	5.23	0.05	5.83	0.04	5.84	0.18	5.84	0.18	5.81	0.27	5.81	0.04	5.75	0.08	5.75	0.27	5.11	0.51	
3 hrs	4.77	0.07	5.09	0.04	5.33	0.22	5.24	0.08	5.24	0.08	5.24	0.29	5.24	0.05	5.11	0.29	5.11	0.29	4.97	0.51	
6 hrs	4.04	0.29	5.24	0.08	4.68	0.10	5.01	0.50	5.01	0.50	5.01	0.51	5.01	0.05	4.97	0.51	4.97	0.51	4.73	0.05	
1 day	3.29	0.60	4.85	0.20	3.89	0.33	4.21	0.08	4.21	0.08	4.21	0.29	4.21	0.05	4.73	0.05	4.73	0.05	4.20	0.07	
2 days	2.47	0.23	4.44	0.20	2.76	0.10	3.16	0.07	3.16	0.07	3.16	0.27	3.16	0.05	3.71	0.50	3.71	0.50	3.21	0.46	
4 days	U	-	3.26	0.10	2.27	0.09	2.47	0.28	2.47	0.28	2.47	0.46	2.47	-	2.79	0.46	2.79	0.46	2.79	0.46	
7 days	U	-	U	-	U	-	U	-	U	-	U	-	U	-	U	-	U	-	U	-	

C) Disinfectants*

D) pH*


Disinfectant (Working concentration)	Virus titre (Log TCID ₅₀ /mL)						pH (60 mins)	Virus titre (Log TCID ₅₀ /mL)		
	5 mins	15 mins	30 mins	Mean	±SD			Mean	±SD	
Household bleach (1:49)	U	U	U	5.55	0.25		3	5.55	0.25	
Household bleach (1:99)	U	U	U	5.67	0.36		4	5.67	0.36	
Hand soap solution (1:49)	3.6 [#]	U	U	5.73	0.04		5	5.73	0.04	
Ethanol (70%)	U	U	U	5.75	0.08		6	5.75	0.08	
Povidone-iodine (7.5%)	U	U	U	5.58	0.22		7	5.58	0.22	
Chloroxylenol (0.05%)	U	U	U	5.70	0.14		8	5.70	0.14	
Chlorhexidine (0.05%)	U	U	U	5.54	0.44		9	5.54	0.44	
Benzalkonium chloride (0.1%)	U	U	U	5.51	0.11		10	5.51	0.11	

* All the virus titres were titrated using Vero-E6 cells. All experimental studies were done in three independent triplicates. Detection limit of a typical TCID₅₀ assay is 100 TCID₅₀/mL, except reactions containing hand soap/chloroxylenol (detection limit: 10³ TCID₅₀/mL) or reactions containing povidone-iodine/cholorhexidin/benzalkonium chloride; detection limit: 10⁴ TCID₅₀/mL because of their cytotoxic effects. N.D.: not done, U: undetectable.

[#] Only one of the triplicate reactions was positive in the TCID₅₀ assay.


Appendix: Biphasic decay of infectious SARS-CoV-2 population

A) Glass




Half-life of free infectious SARS-CoV-2: 0 - 3 hr, $T_{1/2} = 1.2$ hr
 3 hr - 2d, $T_{1/2} = 4.8$ hr

B) Banknote




Half-life of free infectious SARS-CoV-2: 0 - 6 hr, $T_{1/2} = 0.9$ hr
 6 hr - 2d, $T_{1/2} = 7.9$ hr

C) Stainless steel




Half-life of free infectious SARS-CoV-2: 0 -30 min, $T_{1/2} = 0.3$ hr
 30 min - 4d, $T_{1/2} = 14.7$ hr

D) Plastic





Half-life of free infectious SARS-CoV-2: 0 - 6 hr, $T_{1/2} = 1.6$ hr
 6 hr - 4 d, $T_{1/2} = 11.4$ hr

E) Mask, inner layer




Short half-life

Long half-life



F) Mask, outer layer



Half-life of free infectious SARS-CoV-2: 0 - 3 hr, $T_{1/2} = 1.4$ hr
 3 hr - 7 d, $T_{1/2} = 23.9$ hr